


-2-











wherein each hydrogen atom in the radicals (a-1), (a-2), (a-3), (a-4) and (a-5) may optionally be replaced by halo, C₁₋₆alkyl, nitro, amino, hydroxy, C₁₋₆alkyloxy, polyhaloC₁₋₆alkyl, carboxyl, aminoC₁₋₆alkyl, mono- or di(C₁₋₄alkyl)aminoC₁₋₆alkyl, C₁₋₆alkyloxycarbonyl, hydroxyC₁₋₆alkyl, or a radical of formula



wherein $=\text{Z}$ is $=\text{O}$, $=\text{CH}-\text{C}(=\text{O})-\text{NR}^{5a}\text{R}^{5b}$, $=\text{CH}_2$, $=\text{CH}-\text{C}_{1-6}\text{alkyl}$, $=\text{N}-\text{OH}$ or $=\text{N}-\text{O}-\text{C}_{1-6}\text{alkyl}$;

Q is a radical of formula



wherein Alk is C₁₋₆alkanediyl;

Y¹ is a bivalent radical of formula $-\text{NR}^2-$ or $-\text{CH}(\text{NR}^2\text{R}^4)-$;

X¹ is NR⁴, S, S(=O), S(=O)₂, O, CH₂, C(=O), C(=CH₂), CH(OH), CH(CH₃), CH(OCH₃), CH(SCH₃), CH(NR^{5a}R^{5b}), CH₂-NR⁴ or NR⁴-CH₂;

X² is a direct bond, CH₂, C(=O), NR⁴, C₁₋₄alkyl-NR⁴, NR⁴-C₁₋₄alkyl;

t is 2, 3, 4 or 5;

u is 1, 2, 3, 4 or 5;

v is 2 or 3; and

whereby each hydrogen atom in Alk and the carbocycles and the heterocycles defined in radicals (b-3), (b-4), (b-5), (b-6), (b-7) and (b-8) may optionally be replaced

by R^3 ; with the proviso that when R^3 is hydroxy or C_{1-6} alkyloxy, then R^3 can not replace a hydrogen atom in the α position relative to a nitrogen atom;

G is C_{1-10} alkanediyl substituted with one or more hydroxy, C_{1-6} alkyloxy, aryl C_{1-6} alkyloxy, C_{1-6} alkylthio, aryl C_{1-6} alkylthio, $HO(-CH_2-CH_2-O)_n$, C_{1-6} alkyloxy $(-CH_2-CH_2-O)_n$ or aryl C_{1-6} alkyloxy $(-CH_2-CH_2-O)_n$;

R^1 is a monocyclic heterocycle or aryl; said heterocycle being selected from piperidinyl, piperazinyl, pyridyl, pyrazinyl, pyridazinyl, pyrimidinyl, furanyl, tetrahydrofuranyl, thienyl, pyrrolyl, thiazolyl, oxazolyl, imidazolyl, isothiazolyl, pyrazolyl, isoxazolyl, oxadiazolyl; and each heterocycle may optionally be substituted with 1 or where possible more, such as 2, 3 or 4, substituents selected from halo, hydroxy, amino, cyano, carboxy, C_{1-6} alkyl, C_{1-6} alkyloxy, C_{1-6} alkylthio, C_{1-6} alkyloxy C_{1-6} alkyl, aryl, aryl C_{1-6} alkyl, aryl C_{1-6} alkyloxy, hydroxy C_{1-6} alkyl, mono- or di(C_{1-6} alkyl)amino, mono- or di(C_{1-6} alkyl)amino C_{1-6} alkyl, polyhalo C_{1-6} alkyl, C_{1-6} alkylcarbonylamino, C_{1-6} alkyl-SO₂-NR^{5c}-, aryl-SO₂-NR^{5c}-, C_{1-6} alkyloxycarbonyl, -C(=O)-NR^{5c}R^{5d}, $HO(-CH_2-CH_2-O)_n$, halo $(-CH_2-CH_2-O)_n$, C_{1-6} alkyloxy $(-CH_2-CH_2-O)_n$, aryl C_{1-6} alkyloxy $(-CH_2-CH_2-O)_n$ and mono- or di(C_{1-6} alkyl)amino $(-CH_2-CH_2-O)_n$; each n independently is 1, 2, 3 or 4;

R^2 is hydrogen, formyl, C_{1-6} alkylcarbonyl, Hetcarbonyl, pyrrolidinyl, piperidinyl, homopiperidinyl, C_{3-7} cycloalkyl substituted with $N(R^6)_2$, or C_{1-10} alkyl substituted with $N(R^6)_2$ and optionally with a second, third or fourth substituent selected from amino, hydroxy, C_{3-7} cycloalkyl, C_{2-5} alkanediyl, piperidinyl, mono- or di(C_{1-6} alkyl)amino, C_{1-6} alkyloxycarbonylamino, aryl and aryloxy;

R^3 is hydrogen, hydroxy, C_{1-6} alkyl, C_{1-6} alkyloxy, aryl C_{1-6} alkyl or aryl C_{1-6} alkyloxy;

R^4 is hydrogen, C_{1-6} alkyl or aryl C_{1-6} alkyl;

R^{5a} , R^{5b} , R^{5c} and R^{5d} each independently are hydrogen or C_{1-6} alkyl; or

R^{5a} and R^{5b} , or R^{5c} and R^{5d} taken together form a bivalent radical of formula $-(CH_2)_s-$

wherein s is 4 or 5;

R^6 is hydrogen, C_{1-4} alkyl, formyl, hydroxy C_{1-6} alkyl, C_{1-6} alkylcarbonyl or C_{1-6} alkyloxycarbonyl;

contd.

a1

C1
cont 5

10
15
20
25

contd.
a¹

C¹
cont⁵

-4-

aryl is phenyl or phenyl substituted with 1 or more, such as 2, 3 or 4, substituents selected from halo, hydroxy, C₁₋₆alkyl, hydroxyc₁₋₆alkyl, polyhaloC₁₋₆alkyl, and C₁₋₆alkyloxy;

Het is pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl.

2. (amended) A compound according to claim 1, wherein -a¹=a²-a³=a⁴- is a radical of formula (a-1) or (a-2).

10 3. (amended) A compound according to claim 1 wherein R¹ is phenyl optionally substituted with halo, C₁₋₆alkyl or C₁₋₄alkyloxy; or pyridyl optionally substituted with 1 or more substituents selected from arylC₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyl, aryl, mono- or di(C₁₋₆alkyl)amino, C(=O)-NR^{5c}R^{5d}, halo or C₁₋₆alkyl.

15 4. (amended) A compound according to claim 1, wherein G is C₁₋₄alkanediyl substituted with hydroxy, C₁₋₆alkyloxy, HO(-CH₂-CH₂-O)_n-, C₁₋₆alkyloxy(-CH₂-CH₂-O)_n- or arylC₁₋₆alkyloxy(-CH₂-CH₂-O)_n-.

20 5. (amended) A compound according to claim 1, wherein Q is a radical of formula (b-5) wherein v is 2 and Y¹ is -NR²-.

6. (amended) A compound according to claim 1, wherein X¹ is NH or CH₂.

25 7. (amended) A compound according to claim 1, wherein R² is hydrogen or C₁₋₁₀alkyl substituted with NHR⁶ wherein R⁶ is hydrogen or C₁₋₆alkyloxycarbonyl.

30 8. A compound according to claim 1 wherein the compound is [(A),(S)]-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[(6-bromo-2-pyridinyl)ethoxymethyl]-1H-benzimidazol-2-amine; [(A),(S)]-N-[1-(2-aminopropyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine (compound 75); (±)-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[(2-

contd.

a¹

C¹
cont⁵

10093337-0001

-5-

- methoxyethoxy)(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine; N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-6-chloro-1-[(2-methoxyethoxy)(6-methyl-2-pyridinyl)methyl]-4-methyl-1H-benzimidazol-2-amine trihydrochloride trihydrate; [(A),(R)]-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine monohydrate; (±)-N-[1-(2-aminopropyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine; [(A)(S)]-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine monohydrate; (±)-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine; [(A),(R)]-N-[1-(2-aminopropyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine monohydrate; (±)-N-[1-(2-aminoethyl)-4-piperidinyl]-1-[(6-bromo-2-pyridinyl)ethoxymethyl]-2-benzimidazol-2-amine; (±)-N-[1-(2-aminoethyl)-4-piperidinyl]-1-[(2-ethoxyethoxy)(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine; [(B),(S)] N-[1-(2-aminopropyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine monohydrate; (±)-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-3-[(2-methoxyethoxy)(6-methyl-2-pyridinyl)methyl]-7-methyl-3H-imidazo[4,5-b]pyridin-2-amine; (±)-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[(2-ethoxyethoxy)(6-phenyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine; (±)-N-[1-(2-aminoethyl)-4-piperidinyl]-1-[(2-methoxyethoxy)(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine; (±)-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[(6-bromo-2-pyridinyl)ethoxymethyl]-4-methyl-1H-benzimidazol-2-amine monohydrate; [(A),(R)]-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[(6-bromo-2-pyridinyl)ethoxymethyl]-1H-benzimidazol-2-amine; (±)-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[(6-bromo-2-pyridinyl)ethoxymethyl]-1H-benzimidazol-2-amine; a prodrug, N-oxide, addition salt, quaternary amine, metal complex or stereochemically isomeric form thereof.

contd.
a¹

C¹
cont⁵


10
15
20

9. (amended) A method of using as a medicine a compound as claimed in any one of claims 1 to 8.

10. (amended) A pharmaceutical composition, comprising a pharmaceutically acceptable carrier, and as active ingredient a therapeutically effective amount of a compound as claimed in any one of claims 1 to 8.


11. (amended) A process of preparing a composition as claimed in claim 10, comprising the step of intimately mixing said carrier with said compound.

12. An intermediate of formula



with R¹, G and -a¹=a²-a³=a⁴- defined as in claim 1, P being a protective group, and Q₁ being defined as Q according to claim 1 provided that it is devoided of the R² or R⁶ substituent.

13. An intermediate of formula




with R¹, G and -a¹=a²-a³=a⁴- defined as in claim 1, and (O=)Q₃ being a carbonyl derivative of Q, said Q being defined according to claim 1, provided that it is devoided of the -NR²R⁴ or -NR²- substituent.

14. An intermediate of formula

contd.


a¹

C¹
cont



with R¹, Q and -a¹=a²-a³=a⁴- defined as in claim 1, and (O=)G₂ being a carbonyl derivative of G, said G being defined according to claim 1.

- 5 15. (amended) A process of preparing a compound as claimed in claim 1, comprising at least one step selected from the group consisting of:
- a) reacting an intermediate of formula (II-a) or (II-b) with an intermediate of formula (III)



10


with R¹, G, Q and -a¹=a²-a³=a⁴- defined as in claim 1, and W₁ being a suitable leaving group, in the presence of a suitable base and in a suitable reaction-inert solvent;

- b) deprotecting an intermediate of formula (IV)

contd.

a¹


C¹
cont



with R^1 , G , and $-a^1=a^2-a^3=a^4-$ defined as in claim 1, $H-Q_1$ being defined as Q according to claim 1 provided that R^2 or at least one R^6 substituent is hydrogen, and P being a protective group;

5


c) deprotecting and reducing an intermediate of formula (IV-a)



with R^1 , G , and $-a^1=a^2-a^3=a^4-$ defined as in claim 1, $H-Q_1$ being defined as Q according to claim 1 provided that R^2 or at least one R^6 substituent is hydrogen, $Q_{1a}(CH=CH)$ being defined as Q_1 provided that Q_1 comprises an unsaturated bond, and P being a protective group;

10

d) deprotecting an intermediate of formula (V)




with R^1 , G , and $-a^1=a^2-a^3=a^4-$ defined as in claim 1, and H_2N-Q_2 being defined as Q according to claim 1 provided that both R^6 substituents are hydrogen or R^2 and R^4 are both hydrogen;

15

e) deprotecting an intermediate of formula (VI)



contd.
a¹

C1
cont



with R¹, G, and -a¹=a²-a³=a⁴- defined as in claim 1, and H₂N-Q₂ being defined as Q according to claim 1 provided that both R⁶ substituents are hydrogen or R² and R⁴ are both hydrogen, and P being a protective group;

5 f) deprotecting an intermediate of formula (VII) or (VIII)




with R¹, G, and -a¹=a²-a³=a⁴- defined as in claim 1, H-Q₁'(OH) being defined as Q according to claim 1 provided that R² or at least one R⁶ substituent is hydrogen and provided that Q comprises a hydroxy moiety, H₂N-Q₂'(OH) being defined as Q according to claim 1 provided that both R⁶ substituents are hydrogen or R² and R⁴ are both hydrogen and provided that Q comprises a hydroxy moiety, and P being a protective group;

10

g) amination of an intermediate of formula (IX)


contd.
a¹

C¹
cont




with R¹, G, and -a¹=a²-a³=a⁴- defined as in claim 1, and H₂N-Q₃H being defined as Q according to claim 1 provided that both R⁶ substituents are hydrogen or R² and R⁴ are both hydrogen, and the carbon adjacent to the nitrogen carrying the R⁶, or R² and R⁴ substituents contains at least one hydrogen, in the presence of a suitable amination reagent;

h) reducing an intermediate of formula (X)



with R¹, G, and -a¹=a²-a³=a⁴- defined as in claim 1, and H₂N-CH₂-Q₄ being defined as Q according to claim 1 provided that Q comprises a -CH₂-NH₂ moiety, in the presence of a suitable reducing agent;


i) reducing an intermediate of formula (X-a)



with G, and -a¹=a²-a³=a⁴- defined as in claim 1, H₂N-CH₂-Q₄ being defined as Q according to claim 1 provided that Q comprises a -CH₂-NH₂ moiety, and R^{1'} being defined as R¹ according to claim 1 provided that it comprises at least one substituent, in the presence of a suitable reducing agent and suitable solvent;

CI
cont

- THE** **NEW** **YORK** **PUBLIC** **LIBRARY**



- 5




with R^1 , G, and $-a^1=a^2-a^3=a^4$ defined as in claim 1, and R^{2a} -NH-HQ₅ being defined as Q according to claim 1 provided that R^2 is other than hydrogen and is represented by R^{2a} , R^4 is hydrogen, and the carbon atom adjacent to the nitrogen

contd
a¹

C1
contd



atom carrying the R² and R⁴ substituents, carries also at least one hydrogen atom, in the presence of a suitable reducing agent;

m) reducing an intermediate of formula (XV)




5 with R¹, G, and -a¹=a²-a³=a⁴- defined as in claim 1, and (R⁶)₂N-[(C₁-
9alkyl)CH₂OH]-NH-HQ₅ being defined as Q according to claim 1 provided that R²
is other than hydrogen and is represented by C₁₋₁₀alkyl substituted with N(R₆)₂ and
with hydroxy, and the carbon atom carrying the hydroxy, carries also two hydrogen
atoms, and provided that R⁴ is hydrogen, and the carbon atom adjacent to the
10 nitrogen atom carrying the R² and R⁴ substituents, carries also at least one
hydrogen atom, with a suitable reducing agent;

n) deprotecting an intermediate of formula (XVI), (XVI-a) or (XVI-b)




contd
a¹

C¹
cont




with G, and $-a^1=a^2-a^3=a^4-$ defined as in claim 1, and H-Q₁ being defined as Q according to claim 1 provided that R² or at least one R⁶ substituent is hydrogen, and R^{1a}-(A-O-H)_w, R^{1a'}-(A-O-H)₂ and R^{1a''}-(A-O-H)₃ being defined as R¹ according to claim 1 provided that R¹ is substituted with hydroxy, hydroxyC₁₋₆alkyl, or HO(-CH₂-CH₂-O)_n-, with w being an integer from 1 to 4 and P or P₁ being a suitable protecting group, with a suitable acid[.];

o) amination of an intermediate of formula (XVII)




with R¹, G, $-a^1=a^2-a^3=a^4-$, Alk, X¹, R² and R⁴ defined as in claim 1, in the presence of a suitable amination agent;

p) amination of an intermediate of formula (XIX)




with R¹, G, and $-a^1=a^2-a^3=a^4-$ defined as in claim 1, and Q₆N-CH₂-C₁₋₃alkyl-NR⁴ being defined as Q according to claim 1 provided that in the definition of Q, X² is C₂₋₄alkyl-NR⁴, in the presence of a suitable amination agent;

q) deprotecting an intermediate of formula (XXI)



with R^1 , Q , and $-a^1=a^2-a^3=a^4-$ defined as in claim 1, and $HO-G_1$ being defined as G according to claim 1 provided that G is substituted with hydroxy or $HO-(CH_2CH_2O)_n$; and

r) reducing an intermediate of formula (XXII)



with R^1 , Q , and $-a^1=a^2-a^3=a^4-$ defined as in claim 1, and $H-G_2-OH$ being defined as G according to claim 1 provided that G is substituted with hydroxy and the carbon atom carrying the hydroxy substituent carries also at least one hydrogen, in the presence of a suitable reducing agent.

16. (amended) A product, comprising:

(a) a first compound as claimed in claim 1; and

(b) a second antiviral compound, wherein said first compound and said second compound are simultaneously, separately or sequentially used in the treatment or the prevention of viral infections.

17. (amended) A pharmaceutical composition, comprising:

contd.

a¹

C¹
cont

- (a) a pharmaceutically acceptable carrier; and
- (b) as active ingredients:
 - i. a first compound as claimed in claim 1; and
 - ii. a second antiviral compound.

5

Please add the following new claims:

a²

C¹
cont

18
19
20
21

15

20

25

- 18. (new) The process of claim 15, further comprising the step of converting compound of formula (I'), stereochemically isomeric forms, metal complexes, quaternary amines or *N*-oxide forms thereof, into a therapeutically active non-toxic acid addition salt by treatment with an acid.
- 19. (new) The process of claim 15, further comprising the step of converting compound of formula (I'), stereochemically isomeric forms, metal complexes, quaternary amines or *N*-oxide forms thereof, into a therapeutically active non-toxic base addition salt by treatment with alkali.
- 20. (new) The process of claim 15, further comprising the step of converting the acid addition salt form of compound of formula (I'), stereochemically isomeric forms, metal complexes, quaternary amines or *N*-oxide forms thereof, into the free base by treatment with alkali.
- 21. (new) The process of claim 15, further comprising the step of converting the base addition salt form of compound of formula (I'), stereochemically isomeric forms, metal complexes, quaternary amines or *N*-oxide forms thereof, into the free acid by treatment with acid.